arXiv:0905.0273v1 [math.NA] 4 May 2009

Non-globally Lipschitz Counterexamples for the stochastic Euler scheme

Martin Hutzenthaler* and Arnulf Jentzen†

May 4, 2009

Abstract

The stochastic Euler scheme is known to converge to the exact solution of a stochastic differential equation with globally Lipschitz coefficients and even with coefficients which grow at most linearly. For super-linearly growing coefficients convergence in the strong and numerically weak sense remained an open question. In this article we prove for many stochastic differential equations with super-linearly growing coefficients that Euler's approximation does not converge neither in the strong L^p-sense nor in the numerically weak sense to the exact solution. Even worse, the difference of the exact solution and of the numerical approximation diverges to infinity in the strong L^p-sense and in the numerically weak sense.

1 Introduction

An important numerical scheme for simulating stochastic differential equations (SDEs) is Euler's method (see e.g. [14, 20]) (a.k.a. Euler-Maruyama scheme). If the coefficients of an SDE are globally Lipschitz-continuous, then

---

*Research supported by the DFG in the Dutch German Bilateral Research Group "Mathematics of Random Spatial Models from Physics and Biology" (FOR 498)

†Research supported by the DFG project "Pathwise dynamics and numerics of stochastic evolution equations"

AMS subject classifications 65C30.

Euler scheme, stochastic differential equations, counterexamples, global Lipschitz.

standard results (e.g. Chapter 10 and 14 in [14]) show convergence of the Euler approximation in the strong and numerically weak sense to the exact solution of the SDE. It remained an open question whether the Euler approximation also converges in the strong or numerically weak sense if the coefficients are not globally Lipschitz-continuous, see e.g. Section 1 in [8]. In this paper we answer this question to the negative. More precisely we prove for a large class of SDEs with super-linearly growing coefficient functions that both the distance in the strong L^p-sense and the distance between the p-th absolute moments of the Euler approximation and of the exact solution of the SDE diverges to infinity for any $p \in [1, \infty)$. Thus the Euler scheme does in the strong and numerically weak sense not produce an appropriate approximation of the exact solution of such an SDE.

For clarity of exposition we concentrate in this section on the following prominent example (also considered in [19, 21, 23]). Let $(X_t)_{t \geq 0}$ be the unique strong solution of the one-dimensional SDE

$$dX_t = -X_t^3 \, dt + \sigma \, dW_t, \qquad X_0 = x_0, \tag{1}$$

where $(W_t)_{t \geq 0}$ is an one-dimensional standard Brownian motion and $\sigma > 0$, $x_0 \in \mathbb{R}$ are constants. The Euler approximation $(Y_k^N)_{k=0,\dots,N}$ of $(X_t)_{t \in [0,T]}$, $T < \infty$ fixed, is defined recursively through $Y_0^N = x_0$ and

$$Y_{k+1}^N = Y_k^N - \frac{T}{N} \left(Y_k^N \right)^3 + \sigma \left(W_{\frac{(k+1)T}{N}} - W_{\frac{kT}{N}} \right) \tag{2}$$

for every $k = 0, \dots, N-1$ and every $N \in \mathbb{N} := \{1, 2, \dots\}$. We will prove in Section 2 that the L^p-distance of the exact solution X_T and of the numerical approximation Y_N^N diverges to infinity as N tends to infinity, that is,

$$\lim_{N \to \infty} \mathbb{E} \left| X_T - Y_N^N \right|^p = \infty \tag{3}$$

for every $p \in [1, \infty)$, see equation (11). Thus strong convergence of the Euler approximation fails to hold. In addition numerically weak convergence (see e.g. Section 9.4 in [14]) (not to be confused with stochastic weak convergence) fails to hold, that is,

$$\lim_{N \to \infty} \left| \mathbb{E} \left| X_T \right|^p - \mathbb{E} \left| Y_N^N \right|^p \right| = \infty \tag{4}$$

for every $p \in [1, \infty)$, see equation (13). Please note that this divergence of the Euler approximation is not a special property of equation (1). We establish this divergence for a large class of SDEs with super-linearly growing coefficients in Section 3. Moreover our estimates are easily adapted to prove divergence of other numerical schemes such as the Milstein scheme. For simplicity we restrict ourselves to the Euler scheme. Presence of noise, however, is essential. In the deterministic case, e.g. (1) and (2) with $\sigma = 0$, the Euler scheme does converge and both (3) and (4) fail to hold. In addition note that – in contrast to strong and numerically weak convergence – pathwise convergence does hold if the coefficients are locally Lipschitz-continuous, see e.g. Gyöngy (1998), Fleury (2005) or Jentzen et al (2009).

Next we indicate why the L^1-norm of Euler's approximation diverges if the growth of the coefficient functions is not linearly bounded. The Euler approximation of the deterministic equation, that is, (1) and (2) with $\sigma = 0$, is known to be unstable if x_0 is too large (see e.g. Chapter 6 in [3]). A central observation of this article is a quantification of this instability. For example, if $Y_0^N = x_0 = N$ and $T = 1$, then

$$Y_1^N = N - \frac{T}{N} N^3 = N - N^2 \approx -N^2 \tag{5}$$

and therefore

$$Y_2^N = Y_1^N - \frac{T}{N} \left(Y_1^N\right)^3 \approx \left(-N^2\right) - \frac{T}{N} \left(-N^2\right)^3 = -N^2 + N^5 \approx N^5 > N^4$$

and

$$Y_3^N = Y_2^N - \frac{T}{N} \left(Y_2^N\right)^3 \approx \left(N^5\right) - \frac{T}{N} \left(N^5\right)^3 = N^5 - N^{14} \approx -N^{14} < -N^8$$

for every large $N \in \mathbb{N}$. Iterating this calculation, yields $\left|Y_k^N\right| \gtrapprox N^{\left(2^k\right)}$ for every $k = 0, 1, \ldots, N$ and, in particular, $\left|Y_N^N\right| \gtrapprox N^{\left(2^N\right)}$ for every large $N \in \mathbb{N}$. Thus Y_N^N grows double-exponentially fast in N. Now in the presence of noise ($\sigma > 0$) there is an exponentially small event that the Brownian motion leaves the interval $[-2N, 2N]$. On this event the deterministic dynamics lets the approximation grow double-exponentially fast. Consequently as being **double-exponentially large** over-compensates that the event has an **exponentially small** probability, the L^1-norm of the Euler approximation is unbounded in N.

Having sketched why the Euler scheme diverges for many SDEs whose coefficient functions are not linearly bounded we take a short and incomplete look into the literature where we concentrate on the non-globally Lipschitz case and on relatives of the Euler scheme. It is a classical result (e.g. [14, 20]) that the Euler approximation converges in case of globally Lipschitz-continuous functions. In contrast to our results on super-linearly growing coefficients, Yan (2002) proves numerically weak convergence of the Euler scheme if both drift and diffusion function have at most linear growth. Higham et al (2002, Theorem 2.2) provide an instructive condition for strong convergence saying that local Lipschitz-continuity of the coefficients together with boundedness of a p-th moment, $p > 2$, of the approximation and of the exact solution suffices for strong L^2-convergence. Of course our results signify that the p-th moment of the explicit Euler approximation is unbounded if the coefficients grow super-linearly and thus this condition does not apply in that case. Zhang (2006) and Berkaoui et al (2008) prove strong convergence for a class of drift and diffusion functions with a singularity. Yuan and Mao (2008) obtain the rate of convergence of the stochastic Euler scheme for locally Lipschitz-continuous coefficients if these grow at most linearly. An explicit bound for the strong L^1 error is established in Higham and Mao (2005) for the mean-reverting square-root process (a.k.a. CIR process), which has linearly bounded coefficients. Bernard and Fleury (2001) establish convergence in probability under weak hypotheses like local Lipschitz-continuity. Pathwise convergence results can be found in Gyöngy (1998), Fleury (2005), Jentzen and Kloeden (2009) or in Jentzen et al (2009).

A number of authors obtain convergence for modified Euler schemes. Milstein and Tretyakov (2005) consider a modified Euler scheme for non-globally Lipschitz coefficients. They obtain numerically weak convergence by discarding trajectories of the Brownian motion which leave a sufficiently large sphere. As we have seen above, these trajectories are responsible for the divergence of the absolute moments. Lamba et al (2007) prove strong convergence of an Euler scheme with an adaptive time-stepping algorithm for locally Lipschitz-continuous coefficients. A completely different adaptive time-stepping algorithm with focus on long time approximation is proposed by Lemaire (2007). For results on implicit Euler schemes consult e.g. Hu (1996) or also Talay (2002) and the references therein. Finally, Higham et al (2002) showed strong convergence of an implicit Euler scheme via the instructive condition cited above.

Simulations do not always reveal the divergence of the Euler approximation. See Table 1 for simulations in which the approximate value is within distance two of the exact value. To understand this consider the deterministic case, that is, (1) and (2) with $\sigma = 0$. Then the approximation explodes if $|x_0| > \sqrt{2N/T}$. Consequently in the case $\sigma = 1 = T$ it is likely that the approximation diverges if the Brownian path reaches level $\sqrt{2N}$. The probability hereof is bounded above by

$$\mathbb{P}\left[\sup_{0\leq t\leq 1}|W_t|\geq\sqrt{2N}\right]\leq 4\cdot\mathbb{P}\left[W_1\geq\sqrt{2N}\right]\leq\frac{2}{\sqrt{\pi N}}e^{-N}\leq 2e^{-N}\tag{6}$$

for every $N\in\mathbb{N}$, where we have used the reflection principle and a standard estimate, see e.g. Theorem 21.19 in [13]. Thus the probability that we see an explosion of the approximation is negligible for $N\geq 100$, say. Even more if the simulation is repeated N^2 times, say, then the probability to see an explosion is bounded above by about $2N^2e^{-N}$ for every $N\in\mathbb{N}$, which is again negligible for $N\geq 100$, say. Therefore even in a large number of simulations it is unlikely to encounter an approximation which diverges.

The rest of this article is organized as follows. To clearly present the idea of this article, we exemplify the divergence of the Euler scheme for the SDE (1) with $\sigma=1$, $x_0=0$ in Section 2 (Theorem 1). A general result (Theorem 2) and several examples for the divergence of the Euler scheme are provided in Section 3. Simulations in Section 4 illustrate this divergence. The proofs are postponed to the final section.

2 A simple counterexample

Fix $T>0$ and let $(\Omega,\mathcal{F},\mathbb{P})$ be a probability space on which a scalar standard Brownian motion $W\colon[0,T]\times\Omega\to\mathbb{R}$ with continuous sample paths is defined. Then, in this section, we consider the SDE (1) with $\sigma=1$ and $x_0=0$, i.e.

$$dX_t=-X_t^3\,dt+dW_t,\qquad X_0=0\tag{7}$$

for every $t\in[0,T]$. More precisely, let $X\colon[0,T]\times\Omega\to\mathbb{R}$ be the pathwise unique stochastic process with continuous sample paths, that satisfies

$$X_t(\omega)=-\int_0^t(X_s(\omega))^3\,ds+W_t(\omega)\tag{8}$$

for every $t\in[0,T]$ and every $\omega\in\Omega$ (see e.g. Theorem 2.4.1 in [18] for existence and uniqueness of such a process). Then the Euler scheme denoted by $\mathcal{F}/\mathcal{B}(\mathbb{R})$-measurable mappings $Y_k^N\colon\Omega\to\mathbb{R}$, $k=0,1,\ldots,N$, $N\in\mathbb{N}$, is in the setting of equation (7) given by $Y_0^N(\omega)=0$ and

$$Y_{k+1}^N(\omega)=Y_k^N(\omega)-\frac{T}{N}\left(Y_k^N(\omega)\right)^3+\left(W_{\frac{(k+1)T}{N}}(\omega)-W_{\frac{kT}{N}}(\omega)\right)\tag{9}$$

for every $k = 0, 1, \ldots, N-1$, every $N \in \mathbb{N}$ and every $\omega \in \Omega$. We now show the divergence of Euler's method (9) for the SDE (7).

Theorem 1. *Let $T > 0$, the solution X of* (8) *and the Euler approximation Y_k^N, $k = 0, 1, \ldots, N$, $N \in \mathbb{N}$, be as above. Then*

$$\lim_{N \to \infty} \mathbb{E} \left| X_T - Y_N^N \right| = \infty. \tag{10}$$

Remark: Due to Jensen's inequality Theorem 1 in particularly implies

$$\lim_{N \to \infty} \mathbb{E} \left[\left| X_T - Y_N^N \right|^p \right] = \infty \tag{11}$$

for every $p \in [1, \infty)$. In addition since the exact solution $X \colon [0, T] \times \Omega \to \mathbb{R}$ satisfies $\mathbb{E} \left| X_T \right|^p < \infty$ for every $p \in [1, \infty)$ (see e.g. Theorem 2.4.1 in [18]) we get

$$\begin{aligned} \left(\mathbb{E} \left| Y_N^N \right|^p \right)^{\frac{1}{p}} &= \left(\mathbb{E} \left| Y_N^N - X_T + X_T \right|^p \right)^{\frac{1}{p}} \\ &\geq \left(\mathbb{E} \left| Y_N^N - X_T \right|^p \right)^{\frac{1}{p}} - \left(\mathbb{E} \left| X_T \right|^p \right)^{\frac{1}{p}} \longrightarrow \infty \end{aligned} \tag{12}$$

as $N \to \infty$ due to equation (11). Therefore we obtain

$$\lim_{N \to \infty} \left| \mathbb{E} \left| X_T \right|^p - \mathbb{E} \left| Y_N^N \right|^p \right| = \infty \tag{13}$$

for every $p \in [1, \infty)$. $\diamond$

Proof of Theorem 1. First of all we define $r_N := \max\left(3\frac{N}{T}, 1\right)$ for every $N \in \mathbb{N}$. Then we consider the sets

$$\Omega_N := \left\{ \omega \in \Omega \,\middle|\, \sup_{k=1,\ldots,N-1} \left| W_{\frac{(k+1)T}{N}}(\omega) - W_{\frac{kT}{N}}(\omega) \right| \leq 1, \right. \\ \left. \left| W_{\frac{T}{N}}(\omega) - W_0(\omega) \right| \geq r_N \right\}$$

for every $N \in \mathbb{N}$. Note that $\Omega_N \in \mathcal{F}$ for every $N \in \mathbb{N}$.

Let now $N \in \mathbb{N}$ be arbitrary. Then we claim

$$\left| Y_k^N(\omega) \right| \geq (r_N)^{\left(2^{(k-1)}\right)} \qquad \forall \ \omega \in \Omega_N \tag{14}$$

for every $k = 1, 2, \ldots, N$. We prove equation (14) by induction on $k = 1, 2, \ldots, N$. In the base case $k = 1$ we have

$$\left|Y_1^N(\omega)\right| = \left|W_{\frac{T}{N}}(\omega) - W_0(\omega)\right| \geq r_N \tag{15}$$

for every $\omega \in \Omega_N$ due to the definition of the Euler scheme (see equation (9)) and due to the definition of the set Ω_N. Therefore we assume that equation (14) holds for one $k \in \{1, 2, \ldots, N-1\}$. In particular we obtain

$$\left|Y_k^N(\omega)\right| \geq (r_N)^{2^{(k-1)}} \geq r_N \geq 1 \tag{16}$$

for every $\omega \in \Omega_N$. Due to the definition of the set Ω_N, we also have

$$\left|W_{\frac{(k+1)T}{N}}(\omega) - W_{\frac{kT}{N}}(\omega)\right| \leq 1 \tag{17}$$

for every $\omega \in \Omega_N$. Hence, equation (9) and equation (17) imply

$$\begin{aligned}\left|Y_{k+1}^N(\omega)\right| &= \left|Y_k^N(\omega) - \frac{T}{N}\left(Y_k^N(\omega)\right)^3 + \left(W_{\frac{(k+1)T}{N}}(\omega) - W_{\frac{kT}{N}}(\omega)\right)\right| \\ &\geq \left|\frac{T}{N}\left(Y_k^N(\omega)\right)^3\right| - \left|Y_k^N(\omega)\right| - \left|W_{\frac{(k+1)T}{N}}(\omega) - W_{\frac{kT}{N}}(\omega)\right| \\ &\geq \frac{T}{N}\left|Y_k^N(\omega)\right|^3 - \left|Y_k^N(\omega)\right| - 1\end{aligned} \tag{18}$$

for every $\omega \in \Omega_N$. Therefore equation (16) implies

$$\begin{aligned}\left|Y_{k+1}^N(\omega)\right| &\geq \frac{T}{N}\left|Y_k^N(\omega)\right|^3 - 2\left|Y_k^N(\omega)\right|^2 \\ &\geq \left|Y_k^N(\omega)\right|^2\left(\frac{T}{N}\left|Y_k^N(\omega)\right| - 2\right) \\ &\geq \left|Y_k^N(\omega)\right|^2\left(\frac{T}{N}r_N - 2\right) \geq \left|Y_k^N(\omega)\right|^2\end{aligned} \tag{19}$$

for every $\omega \in \Omega_N$, since $\frac{T}{N} r_N - 2 \geq 1$ by definition. Therefore the induction hypothesis, i.e. that equation (14) holds for k, implies

$$\left|Y_{k+1}^N(\omega)\right| \geq \left((r_N)^{(2^{k-1})}\right)^2 = (r_N)^{(2^k)} \tag{20}$$

for all $\omega \in \Omega_N$, which indeed shows that equation (14) holds for $k+1$. Hence, we have shown equation (14) for every $k = 1, 2, \ldots, N$ and since $N \in \mathbb{N}$ was arbitrary, we obtain

$$\left|Y_N^N(\omega)\right| \geq (r_N)^{\left(2^{(N-1)}\right)} \tag{21}$$

for every $\omega \in \Omega_N$ and every $N \in \mathbb{N}$. This yields

$$\mathbb{E}\left|Y_N^N\right| = \int_\Omega \left|Y_N^N(\omega)\right| \mathbb{P}(d\omega) \geq \int_{\Omega_N} \left|Y_N^N(\omega)\right| \mathbb{P}(d\omega) \geq \mathbb{P}\left[\Omega_N\right] \cdot 2^{\left(2^{(N-1)}\right)} \tag{22}$$

for every $N \in \mathbb{N}$. Since

$$\begin{aligned}
\mathbb{P}\left[\Omega_N\right] &= \mathbb{P}\left[\sup_{k=1,\ldots,N-1} \left|W_{\frac{(k+1)T}{N}} - W_{\frac{kT}{N}}\right| \leq 1\right] \cdot \mathbb{P}\left[\left|W_{\frac{T}{N}}\right| \geq r_N\right] \\
&\geq \mathbb{P}\left[\sup_{0\leq t\leq T} |W_t| \leq \frac{1}{2}\right] \cdot \mathbb{P}\left[\left|W_{\frac{T}{N}}\right| \geq r_N\right] \\
&= \mathbb{P}\left[\sup_{0\leq t\leq T} |W_t| \leq \frac{1}{2}\right] \cdot \mathbb{P}\left[\sqrt{\frac{N}{T}}\left|W_{\frac{T}{N}}\right| \geq \sqrt{\frac{N}{T}} r_N\right] \\
&\geq \mathbb{P}\left[\sup_{0\leq t\leq T} |W_t| \leq \frac{1}{2}\right] \cdot \frac{1}{4}\left(\sqrt{\frac{N}{T}} r_N\right) e^{-\left(\sqrt{\frac{N}{T}} r_N\right)^2} \\
&\geq \frac{1}{4\sqrt{T}} \cdot \mathbb{P}\left[\sup_{0\leq t\leq T} |W_t| \leq \frac{1}{2}\right] \cdot e^{-\frac{N}{T}(r_N)^2}
\end{aligned} \tag{23}$$

for every $N \in \mathbb{N}$ due to Lemma 1 in Section 5 below, we obtain

$$\mathbb{E}\left|Y_N^N\right| \geq \frac{1}{4\sqrt{T}} \cdot \mathbb{P}\left[\sup_{0\leq t\leq T} |W_t| \leq \frac{1}{2}\right] \cdot e^{-\frac{N}{T}(r_N)^2} \cdot 2^{\left(2^{(N-1)}\right)} \tag{24}$$

for every $N \in \mathbb{N}$. This shows

$$\begin{aligned}
\lim_{N\to\infty} \mathbb{E}\left|Y_N^N\right| &\geq \frac{1}{4\sqrt{T}} \cdot \mathbb{P}\left[\sup_{0\leq t\leq T} |W_t| \leq \frac{1}{2}\right] \cdot \lim_{N\to\infty}\left(e^{-\frac{N}{T}(r_N)^2} \cdot 2^{\left(2^{(N-1)}\right)}\right) \\
&= \frac{1}{4\sqrt{T}} \cdot \mathbb{P}\left[\sup_{0\leq t\leq T} |W_t| \leq \frac{1}{2}\right] \cdot \lim_{N\to\infty}\left(e^{-\left(9\frac{N^3}{T^3}\right)} \cdot 2^{\left(2^{(N-1)}\right)}\right) = \infty.
\end{aligned}$$

Since, as mentioned above, the exact solution $X : [0, T] \times \Omega \to \mathbb{R}$ of the SDE (7) satisfies $\mathbb{E}|X_T| < \infty$ (see e.g. Theorem 2.4.1 in [18]), we obtain

$$\lim_{N\to\infty} \mathbb{E}\left|X_T - Y_N^N\right| \geq \lim_{N\to\infty} \mathbb{E}\left|Y_N^N\right| - \mathbb{E}|X_T| = \infty, \tag{25}$$

which is the assertion. □

3 Further counterexamples

Throughout this section assume the following setting. Fix $T > 0$ and let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space with normal filtration $(\mathcal{F}_t)_{t\in[0,T]}$. In addition let $W\colon [0,T]\times\Omega \to \mathbb{R}$ be a scalar standard Brownian motion with respect to $(\mathcal{F}_t)_{t\in[0,T]}$ with continuous sample paths and let $\xi\colon \Omega \to \mathbb{R}$ be a $\mathcal{F}_0/\mathcal{B}(\mathbb{R})$-measurable mapping. Let $f, g\colon \mathbb{R} \to \mathbb{R}$ be two $\mathcal{B}(\mathbb{R})/\mathcal{B}(\mathbb{R})$-measurable functions such that the SDE

$$dX_t = f(X_t)\,dt + g(X_t)\,dW_t, \qquad X_0 = \xi, \qquad t \in [0,T] \tag{26}$$

has a solution $X\colon [0,T]\times\Omega \to \mathbb{R}$. More precisely we assume that $X\colon [0,T]\times\Omega \to \mathbb{R}$ is a predictable stochastic process satisfying

$$\mathbb{P}\left[\int_0^T |f(X_s)| + |g(X_s)|^2\,ds < \infty\right] = 1 \tag{27}$$

and

$$\mathbb{P}\left[X_t = \xi + \int_0^t f(X_s)\,ds + \int_0^t g(X_s)\,dW_s\right] = 1 \tag{28}$$

for every $t \in [0,T]$. Then Euler's approximation for the SDE (26) – denoted by $\mathcal{F}/\mathcal{B}(\mathbb{R})$-measurable mappings $Y_k^N\colon \Omega \to \mathbb{R}$, $k = 0, 1, \ldots, N$, $N \in \mathbb{N}$ – is given by $Y_0^N(\omega) = \xi(\omega)$ and

$$Y_{k+1}^N(\omega) = Y_k^N(\omega) + \frac{T}{N} f\left(Y_k^N(\omega)\right) + g\left(Y_k^N(\omega)\right)\left(W_{\frac{(k+1)T}{N}}(\omega) - W_{\frac{kT}{N}}(\omega)\right) \tag{29}$$

for every $k = 0, 1, \ldots, N-1$, every $N \in \mathbb{N}$ and every $\omega \in \Omega$. Now we formalise the main result of this article which asserts the divergence of Euler's method for the SDE (26) with super-linearly growing coefficients.

Theorem 2. *Assume that the setting above is fulfilled with* $\mathbb{P}\left[\,g(\xi) \neq 0\,\right] > 0$ *and let* $C \geq 1$, $\beta > \alpha > 1$ *be constants such that*

$$\max\left(|f(x)|, |g(x)|\right) \geq \frac{|x|^\beta}{C}, \qquad \min\left(|f(x)|, |g(x)|\right) \leq C\,|x|^\alpha \tag{30}$$

for all $|x| \geq C$. *If the exact solution of* (26) *satisfies* $\mathbb{E}\,|X_T|^p < \infty$ *for one* $p \in [1, \infty)$, *then*

$$\lim_{N\to\infty} \mathbb{E}\left|X_T - Y_N^N\right|^p = \infty \qquad \text{and} \qquad \lim_{N\to\infty} \left|\mathbb{E}\,|X_T|^p - \left|Y_N^N\right|^p\right| = \infty, \tag{31}$$

where Y_k^N, $k = 0, 1, \ldots, N$, $N \in \mathbb{N}$, *is defined in equation* (29).

Condition (30) should be read as follows. Either the drift function or the diffusion function grows in a higher order than linearly and the remaining function grows slower than that. More formally it suffices to show that either

$$|f(x)| \geq \frac{|x|^{\beta}}{C}, \qquad |g(x)| \leq C\,|x|^{\alpha} \tag{32}$$

for all $|x| \geq C$ or

$$|g(x)| \geq \frac{|x|^{\beta}}{C}, \qquad |f(x)| \leq C\,|x|^{\alpha} \tag{33}$$

for all $|x| \geq C$ and for some constants $\beta > 1$, $\beta > \alpha \geq 0$, $C > 0$. Please note that our estimates need $\beta > 1$. A super-linear drift function such as $f(x) = x \log x$ is too small for our estimates. The assumption that the diffusion function does not vanish on the starting point ensures presence of noise in the first time step.

The proof of Theorem 2 is deferred to Section 5. In the remainder of this section we apply Theorem 2 to a selection of examples. Please note that the coefficients in the following examples are locally Lipschitz-continuous and satisfy an appropriate one-sided Lipschitz condition. Therefore the p-th absolute moment in each example is finite for every $p \in [1, \infty)$ according to Theorem 2.4.1 in [18]. For every example we check that assumption (30) is satisfied. Thus both the distance in the strong L^p-sense and the distance between the p-th absolute moments of the Euler approximation and of the exact solution diverges to infinity for any $p \in [1, \infty)$ in each of the following examples.

The introductory example The example in Section 1 is

$$dX_t = -X_t^3 dt + \sigma dW_t, \qquad X_0 = x_0 \in \mathbb{R} \tag{34}$$

where $\sigma > 0$. The dominating coefficient is the drift function with dominating exponent $\beta = 3$. The diffusion function has exponent zero and we may choose $\alpha = 0$. The constant C in equation (32) can be chosen as $C = \max(1, \sigma)$.

Stochastic Ginzburg-Landau equation The Ginzburg-Landau equation is from the theory of superconductivity. It has been introduced by

Ginzburg and Landau (1950) [5] to describe a phase transition. Its stochastic version with multiplicative noise can be written as

$$dX_t = \left((\eta + \frac{1}{2}\sigma^2)X_t - \lambda X_t^3\right) dt + \sigma X_t dW_t, \qquad X_0 = x_0 \in (0, \infty) \tag{35}$$

where $\eta \geq 0$, $\lambda, \sigma > 0$. Its solution is known explicitly (see e.g. Section 4.4 in [14])

$$X_t = \frac{x_0 \exp\left(\eta t + \sigma W_t\right)}{\sqrt{1 + 2x_0^2 \lambda \int_0^t \exp\left(2\eta s + 2\sigma W_s\right) ds}} \tag{36}$$

for $t \geq 0$. Since

$$\begin{aligned} \left|\left(\eta + \frac{\sigma^2}{2}\right) x - \lambda x^3\right| &\geq \lambda \left|x\right|^3 - \left(\eta + \frac{\sigma^2}{2}\right) |x| \\ &\geq \frac{\lambda}{2} \left|x\right|^3 + \frac{\lambda}{2} \left|x\right|^3 - \left(\eta + \frac{\sigma^2}{2}\right) |x| \\ &= \frac{\lambda}{2} \left|x\right|^3 + \frac{\lambda}{2} |x| \left(\left|x\right|^2 - \frac{2\eta + \sigma^2}{\lambda}\right) \\ &\geq \frac{\lambda}{2} \left|x\right|^3 + \frac{\lambda}{2} |x| \left(|x| - \frac{2\eta + \sigma^2}{\lambda}\right) \geq \frac{\lambda}{2} \left|x\right|^3 \end{aligned} \tag{37}$$

for all $|x| \geq \max\left(1, \frac{2\eta+\sigma^2}{\lambda}\right)$, the constants in equation (32) can be chosen as $\beta = 3$, $\alpha = 1$, $C = \max\left(1, \sigma, \frac{2}{\lambda}, \frac{2\eta+\sigma^2}{\lambda}\right)$.

Stochastic Verhulst equation The Verhulst equation is an ordinary differential equation and is a simple model for a population with competition between individuals. Its stochastic version with multiplicative noise can be written as

$$dX_t = \left((\eta + \frac{1}{2}\sigma^2)X_t - \lambda X_t^2\right) dt + \sigma X_t dW_t, \qquad X_0 = x_0 \in (0, \infty) \tag{38}$$

where $\eta, \lambda, \sigma > 0$. Its solution is known explicitly (see e.g. Section 4.4 in [14]) and is given by

$$X_t = \frac{x_0 \exp\left(\eta t + \sigma W_t\right)}{1 + x_0 \lambda \int_0^t \exp\left(\eta s + \sigma W_s\right) ds} \tag{39}$$

for $t \geq 0$. The dominant exponent of the drift function is two and of the diffusion function is one. Thus we may choose $\beta = 2$ and $\alpha = 1$. The constant C in equation (32) can be chosen as $C = \max\left(\sigma, \frac{2}{\lambda}, \frac{2\eta+\sigma^2}{\lambda}\right)$.

Feller diffusion with logistic growth The branching process with logistic growth (see e.g. [16]) is a stochastic Verhulst equation with Feller noise. It solves

$$dX_t = \lambda X_t \left(K - X_t\right) dt + \sigma \sqrt{X_t}\, dW_t, \qquad X_0 = x_0 \in (0, \infty) \tag{40}$$

where $\lambda, K, \sigma > 0$. There is no explicit solution for this equation. However it features the following self-duality

$$\mathbb{E}^x \exp\left(-\frac{2\lambda}{\sigma^2} X_t y\right) = \mathbb{E}^y \exp\left(-\frac{2\lambda}{\sigma^2} x X_t\right), \qquad \forall\, t, x, y \geq 0, \tag{41}$$

where $\mathbb{E}^x$ refers to the starting point $X_0 = x$, see [10]. As constants for equation (32) serve $\beta = 2$, $\alpha = 1$, $C = \max(1, \frac{2}{\lambda}, \sigma, 2K)$.

Ohta-Kimura model The Ohta-Kimura model ([22]) is a population model with two types, fixed total population and random selection. Its diffusion equation is

$$dX_t = \sigma X_t (1 - X_t)\, dW_t, \qquad X_0 = x_0 \in (0, 1) \tag{42}$$

where $\sigma > 0$. Since

$$\begin{aligned} |\sigma x(1-x)| &\geq \sigma |x|^2 - \sigma |x| = \frac{\sigma}{2} |x|^2 + \frac{\sigma}{2} |x|^2 - \sigma |x| \\ &= \frac{\sigma}{2} |x|^2 + \frac{\sigma}{2} |x| \left(|x| - 2\right) \geq \frac{\sigma}{2} |x|^2 \end{aligned} \tag{43}$$

for all $|x| \geq 2$, Theorem 2 applies with $\beta = 2$, $\alpha = 0$ and $C = \max(2, \frac{2}{\sigma})$ in equation (33). Note that this is an example with the diffusion function being the dominant coefficient.

4 Simulations

In this section we present simulations which illustrate the dichotomy between the double-exponential growth of the deterministic dynamics (for some initial values) and the exponentially small probability of the event that the noise is sufficiently large. For this purpose we choose an equation with an explicit solution to compare with. Consider the Stratonovich equation

$$dX_t = -X_t^3\, dt + \sigma X_t \circ dW_t, \qquad X_0 = 1, \qquad t \in [0, 3]. \tag{44}$$

This equation agrees with the stochastic Ginzburg-Landau equation (35) in the case $\eta = 0, \lambda = 1, x_0 = 1$. Its explicit solution is given by equation (36) with $\eta = 0, \lambda = 1, x_0 = 1$.

Table 1 shows Monte Carlo simulations of the second moment both of the explicit solution and of the Euler approximation for five values of σ. For each value of σ the table contains ten simulation runs. The number of time steps is $N = 10^3$. In the case $\sigma = 7$ the parameters of the simulation are such that the Euler approximation produces the value "NaN" (NaN is the IEEE arithmetic representation for "not-a-number", here this is due to an operation "Inf $-$ Inf" where Inf is the IEEE arithmetic representation for positive infinity). In contrast to this the Monte Carlo simulations in the cases $\sigma \in \{2, 4, 5\}$ are all finite. As explained in the introduction the Euler approximation grows double-exponentially fast only on an event of an exponentially small probability. For $\sigma \in \{2, 4, 5\}$ this probability is so small that none of the simulation runs produced a large trajectory. The last but one column in Table 1 exemplifies a parameter setting in which the event of large growth has a probability such that in some Monte Carlo simulations no explosion occurs and in some Monte Carlo simulations at least one explosion in 10^5 runs occurs.

	$\sigma = 2$	$\sigma = 4$	$\sigma = 5$	$\sigma = 6$	$\sigma = 7$
Exact solution $\mathbb{E}\left[(X_3)^2\right]$	0.4739	0.9598	1.2357	1.5423	1.8861
Simulation 1	0.4577	0.7448	0.7243	NaN	NaN
Simulation 2	0.4549	0.7518	0.7094	0.5097	NaN
Simulation 3	0.4632	0.7330	0.6912	NaN	NaN
Simulation 4	0.4580	0.7544	0.7150	0.5378	NaN
Simulation 5	0.4619	0.7458	0.6998	0.5197	NaN
Simulation 6	0.4610	0.7664	0.7327	0.5243	NaN
Simulation 7	0.4606	0.7542	0.7189	NaN	NaN
Simulation 8	0.4690	0.7451	0.7122	NaN	NaN
Simulation 9	0.4667	0.7336	0.7146	0.5475	NaN
Simulation 10	0.4591	0.7605	0.6659	NaN	NaN

Table 1: Simulation for the SDE (44). The number of time steps is $N = 10^3$. The number of Monte Carlo runs for the approximation is 10^5 and for the exact solution 10^7.

The values in Table 1 are either within radius two of the true value or

"NaN". This is due to the double-exponential growth of the deterministic system for some initial values. If the simulation starts to grow, then it reaches "NaN" very quickly. We encountered similar behaviour for other exponents greater than two. In order to see double-exponential growth of the Euler approximation in a plot we consider an exponent close to one. More precisely we plot the Monte Carlo simulation of the first absolute moment of the Euler approximation of X_{10} where



Figure 1: First absolute moment $\mathbb{E}\left|Y_N^N\right|$ of the Euler scheme for number of time steps $N = 1, 2, 3, \ldots, 50$. Number of Monte Carlo runs: 10.000

$$dX_t = -10\,\mathrm{sgn}(X_t)\,|X_t|^{1.1}\,dt + 4\,dW_t, \qquad X_0 = 0, \tag{45}$$

see Figure 1 (see also Table 2). Please note that the graph resembles an exponential function and that the y-axis is logarithmic. Thus the growth of the graph in Figure 1 is indeed close to a double-exponential function. In addition we should mention that some fine-tuning was necessary to obtain

N	$\mathbb{E}\left\|Y_N^N\right\|$
$N=1$	10.1160
$N=2$	444.5797
$N=3$	$1.3649 \cdot 10^4$
$N=4$	$3.7566 \cdot 10^5$
$N=5$	$1.0295 \cdot 10^7$
$\vdots$	$\vdots$

N	$\mathbb{E}\left\|Y_N^N\right\|$
$N=48$	$3.4449 \cdot 10^{260}$
$N=49$	$2.6631 \cdot 10^{271}$
$N=50$	$6.1140 \cdot 10^{297}$
$N=51$	$7.2342 \cdot 10^{301}$
$N=52$	Inf
$N=53$	Inf

Table 2: Simulation values of the first absolute moment $\mathbb{E}\left|Y_N^N\right|$ for $N = 1, 2, 3, \ldots, 53$

suitable parameters for which the simulated absolute moment grows but is not "NaN".

5 Proofs

Lemma 1. *Let* $(\Omega, \mathcal{F}, \mathbb{P})$ *be a probability space and let* $Z\colon \Omega \to \mathbb{R}$ *be an* $\mathcal{F}/\mathcal{B}(\mathbb{R})$*-measurable mapping which is standard normal distributed. Then we have*

$$\mathbb{P}\Big[\, |Z| \geq x \,\Big] \geq \frac{x\, e^{-x^2}}{4}, \qquad \mathbb{P}\Big[\, |Z| \in [x, 2x] \,\Big] \geq \frac{x\, e^{-2x^2}}{2} \tag{46}$$

for every $x \in [0, \infty)$.

Proof of Lemma 1. We have

$$\begin{aligned} \mathbb{P}\left[\, |Z| \geq x \,\right] &= 2 \cdot \mathbb{P}\left[\, Z \geq x \,\right] = 2 \left(\int_x^\infty \frac{1}{\sqrt{2\pi}} e^{-\frac{s^2}{2}} ds \right) \\ &\geq 2 \left(\int_x^{\sqrt{2}x} \frac{1}{\sqrt{2\pi}} e^{-\frac{s^2}{2}} ds \right) \\ &\geq 2 \left(\sqrt{2} - 1 \right) x \left(\frac{1}{\sqrt{2\pi}} e^{-\frac{(\sqrt{2}x)^2}{2}} \right) \\ &= \left(\frac{2\left(\sqrt{2}-1\right)}{\sqrt{2\pi}} \right) x e^{-x^2} \geq \frac{x e^{-x^2}}{4} \end{aligned} \tag{47}$$

for every $x \in [0, \infty)$. Moreover, we have

$$\begin{aligned} \mathbb{P}\left[\,|Z| \in [x, 2x]\,\right] &= 2 \cdot \mathbb{P}\left[\,Z \in [x, 2x]\,\right] = 2\left(\int_x^{2x} \frac{1}{\sqrt{2\pi}} e^{-\frac{s^2}{2}} ds\right) \\ &\geq 2x\left(\frac{1}{\sqrt{2\pi}} e^{-\frac{(2x)^2}{2}}\right) = \left(\frac{2}{\sqrt{2\pi}}\right) x e^{-2x^2} \geq \frac{x\, e^{-2x^2}}{2} \end{aligned} \tag{48}$$

for every $x \in [0, \infty)$. □

Proof of Theorem 2. First of all we have $\mathbb{P}[\,|g(\xi)| > 0\,] > 0$. Therefore there exists a real number $K > 1$, such that

$$\mu := \mathbb{P}\left[\,|g(\xi)| \geq \frac{1}{K},\;\; |\xi| + T\,|f(\xi)| \leq K\right] > 0. \tag{49}$$

Then we define

$$r_N := \max\left(2, C, \left(\frac{2CN}{T} + 2C^2\right)^{\frac{1}{(\beta-\alpha)}}\right) \in [C, \infty) \tag{50}$$

and consider the sets $\Omega_N \in \mathcal{F}$ given by

$$\begin{aligned} \Omega_N := \Bigg\{\omega \in \Omega \,\Bigg|\, & \left|W_{\frac{(k+1)T}{N}}(\omega) - W_{\frac{kT}{N}}(\omega)\right| \in \left[\frac{T}{N}, \frac{2T}{N}\right] \forall\, k = 1, \ldots, N-1, \\ & \left|W_{\frac{T}{N}}(\omega) - W_0(\omega)\right| \geq K\,(r_N + K)\,, \\ & |g(\xi(\omega))| \geq \frac{1}{K},\;\; |\xi(\omega)| + T\,|f(\xi(\omega))| \leq K\Bigg\} \end{aligned} \tag{51}$$

for every $N \in \mathbb{N}$. Note that

$$\frac{T}{NC}\,(r_N)^{(\beta-\alpha)} \geq \frac{T}{NC}\left(\frac{2CN}{T} + 2C^2\right) \geq 2 + \frac{2TC}{N} \tag{52}$$

for every $N \in \mathbb{N}$ due to the definition of r_N, $N \in \mathbb{N}$.

Let now $N \in \mathbb{N}$ be arbitrary. Then we claim

$$\left|Y_k^N(\omega)\right| \geq (r_N)^{\left(\alpha^{(k-1)}\right)} \qquad \forall\;\, \omega \in \Omega_N \tag{53}$$

for every $k = 1, 2, \ldots, N$. We prove equation (53) by induction on $k \in \{1, 2, \ldots, N\}$. In the base case $k = 1$ we have

$$\begin{aligned}\left|Y_1^N(\omega)\right| &= \left|\xi(\omega) + \frac{T}{N} f\left(\xi(\omega)\right) + g\left(\xi(\omega)\right)\left(W_{\frac{T}{N}}(\omega) - W_0(\omega)\right)\right| \\ &\geq \left|g\left(\xi(\omega)\right)\right| \cdot \left|W_{\frac{T}{N}}(\omega) - W_0(\omega)\right| - |\xi(\omega)| - \frac{T}{N}\left|f\left(\xi(\omega)\right)\right| \\ &\geq \left|g\left(\xi(\omega)\right)\right| \cdot \left|W_{\frac{T}{N}}(\omega) - W_0(\omega)\right| - K \\ &\geq \frac{1}{K}\left|W_{\frac{T}{N}}(\omega) - W_0(\omega)\right| - K \geq \frac{K\left(r_N + K\right)}{K} - K = r_N\end{aligned} \tag{54}$$

for every $\omega \in \Omega_N$ due to equation (29) and equation (51). For the induction step $k \to k+1$ we assume that equation (53) holds for one $k \in \{1, 2, \ldots, N-1\}$. In particular we obtain

$$\left|Y_k^N(\omega)\right| \geq r_N \geq C \tag{55}$$

for every $\omega \in \Omega_N$, since $r_N \geq 2$. We also have

$$\begin{aligned}\left|Y_{k+1}^N(\omega)\right| &= \left|Y_k^N(\omega) + \frac{T}{N} f(Y_{k+1}^N(\omega)) + g(Y_{k+1}^N(\omega))\left(W_{\frac{(k+1)T}{N}}(\omega) - W_{\frac{kT}{N}}(\omega)\right)\right| \\ &\geq \left|\frac{T}{N} f(Y_k^N(\omega)) + g(Y_k^N(\omega))\left(W_{\frac{(k+1)T}{N}}(\omega) - W_{\frac{kT}{N}}(\omega)\right)\right| - \left|Y_k^N(\omega)\right| \\ &\geq \max\left(\frac{T}{N}\left|f(Y_k^N(\omega))\right|, \left|g(Y_k^N(\omega))\right| \cdot \left|W_{\frac{(k+1)T}{N}}(\omega) - W_{\frac{kT}{N}}(\omega)\right|\right) \\ &\quad - \min\left(\frac{T}{N}\left|f(Y_k^N(\omega))\right|, \left|g(Y_k^N(\omega))\right| \cdot \left|W_{\frac{(k+1)T}{N}}(\omega) - W_{\frac{kT}{N}}(\omega)\right|\right) \\ &\quad - \left|Y_k^N(\omega)\right|\end{aligned}$$

and

$$\begin{aligned}\left|Y_{k+1}^N(\omega)\right| &\geq \max\left(\frac{T}{N}\left|f(Y_k^N(\omega))\right|, \frac{T}{N}\left|g(Y_k^N(\omega))\right|\right) \\ &\quad - \min\left(\frac{T}{N}\left|f(Y_k^N(\omega))\right|, \frac{2T}{N}\left|g(Y_k^N(\omega))\right|\right) - \left|Y_k^N(\omega)\right| \\ &\geq \frac{T}{N}\max\left(\left|f(Y_k^N(\omega))\right|, \left|g(Y_k^N(\omega))\right|\right) \\ &\quad - \frac{2T}{N}\min\left(\left|f(Y_k^N(\omega))\right|, \left|g(Y_k^N(\omega))\right|\right) - \left|Y_k^N(\omega)\right|\end{aligned} \tag{56}$$

for every $\omega \in \Omega_N$ due to equation (29) and equation (51). Therefore equation (55) and equation (30) yield

$$\begin{aligned} \left|Y_{k+1}^N(\omega)\right| &\geq \frac{T}{NC}\left|Y_k^N(\omega)\right|^{\beta} - \frac{2TC}{N}\left|Y_k^N(\omega)\right|^{\alpha} - \left|Y_k^N(\omega)\right| \\ &\geq \frac{T}{NC}\left|Y_k^N(\omega)\right|^{\beta} - \frac{2TC}{N}\left|Y_k^N(\omega)\right|^{\alpha} - \left|Y_k^N(\omega)\right|^{\alpha} \\ &= \left|Y_k^N(\omega)\right|^{\alpha}\left(\frac{T}{NC}\left|Y_k^N(\omega)\right|^{(\beta-\alpha)} - \frac{2TC}{N} - 1\right) \\ &\geq \left|Y_k^N(\omega)\right|^{\alpha}\left(\frac{T}{NC}\left(r_N\right)^{(\beta-\alpha)} - \frac{2TC}{N} - 1\right) \geq \left|Y_k^N(\omega)\right|^{\alpha} \end{aligned} \tag{57}$$

for every $\omega \in \Omega_N$ where the last step follows from equation (52). Hence the induction hypothesis yields

$$\left|Y_{k+1}^N(\omega)\right| \geq \left|Y_k^N(\omega)\right|^{\alpha} \geq \left(\left(r_N\right)^{\left(\alpha^{(k-1)}\right)}\right)^{\alpha} = \left(r_N\right)^{\left(\alpha^{k}\right)} \tag{58}$$

for every $\omega \in \Omega_N$, which proves equation (53) for $k+1$. Therefore equation (53) indeed holds for every $k \in \{1, 2, \ldots, N\}$.

In particular – since $N \in \mathbb{N}$ was arbitrary – we obtain

$$\left|Y_N^N(\omega)\right| \geq \left(r_N\right)^{\left(\alpha^{(N-1)}\right)} \tag{59}$$

for every $\omega \in \Omega_N$ and every $N \in \mathbb{N}$. Thus

$$\begin{aligned} \mathbb{E}\left|Y_N^N\right| &= \int_{\Omega}\left|Y_N^N(\omega)\right| \mathbb{P}(d\omega) \geq \int_{\Omega_N}\left|Y_N^N(\omega)\right| \mathbb{P}(d\omega) \\ &\geq \int_{\Omega_N}\left(r_N\right)^{\left(\alpha^{(N-1)}\right)} \mathbb{P}(d\omega) = \mathbb{P}\left[\Omega_N\right] \cdot \left(r_N\right)^{\left(\alpha^{(N-1)}\right)} \geq \mathbb{P}\left[\Omega_N\right] \cdot 2^{\left(\alpha^{(N-1)}\right)} \end{aligned}$$

for every $N \in \mathbb{N}$. Hence we obtain

$$\begin{aligned}
\mathbb{E}\left|Y_N^N\right| &\geq \mathbb{P}\left[\left|W_{\frac{(k+1)T}{N}} - W_{\frac{kT}{N}}\right| \in \left[\frac{T}{N}, \frac{2T}{N}\right] \forall\, k = 1, \ldots, N-1\right] \\
&\quad \cdot \mathbb{P}\left[\left|W_{\frac{T}{N}} - W_0\right| \geq K\left(r_N + K\right)\right] \cdot \mu \cdot 2^{\left(\alpha^{(N-1)}\right)} \\
&= \left(\mathbb{P}\left[\left|W_{\frac{T}{N}}\right| \in \left[\frac{T}{N}, \frac{2T}{N}\right]\right]\right)^{(N-1)} \cdot \mathbb{P}\left[\left|W_{\frac{T}{N}}\right| \geq K\left(r_N + K\right)\right] \\
&\quad \cdot \mu \cdot 2^{\left(\alpha^{(N-1)}\right)} \\
&\geq \left(\mathbb{P}\left[\sqrt{\frac{N}{T}}\left|W_{\frac{T}{N}}\right| \in \left[\frac{\sqrt{T}}{\sqrt{N}}, \frac{2\sqrt{T}}{\sqrt{N}}\right]\right]\right)^{N} \\
&\quad \cdot \mathbb{P}\left[\sqrt{\frac{N}{T}}\left|W_{\frac{T}{N}}\right| \geq \sqrt{\frac{N}{T}} K\left(r_N + K\right)\right] \cdot \mu \cdot 2^{\left(\alpha^{(N-1)}\right)}
\end{aligned} \tag{60}$$

for every $N \in \mathbb{N}$, where $\mu > 0$ is defined in (49). Hence, Lemma 1 yields

$$\begin{aligned}
\mathbb{E}\left|Y_N^N\right| &\geq \left(\frac{1}{2}\sqrt{\frac{T}{N}} e^{-2\frac{T}{N}}\right)^{N} \cdot \left(\sqrt{\frac{N}{T}} K\left(r_N + K\right)\right) \cdot e^{-\left(\sqrt{\frac{N}{T}} K\left(r_N + K\right)\right)^2} \\
&\quad \cdot \frac{\mu}{4} \cdot 2^{\left(\alpha^{(N-1)}\right)} \\
&\geq \left(\frac{1}{2}\sqrt{\frac{T}{N}}\right)^{N} \cdot e^{-\left(\sqrt{\frac{N}{T}} K\left(r_N + K\right)\right)^2} \cdot \frac{\mu\, e^{-2T}}{4\sqrt{T}} \cdot 2^{\left(\alpha^{(N-1)}\right)} \\
&\geq \left(\frac{1}{2}\sqrt{\frac{T}{N}}\right)^{N} \cdot e^{-\frac{N}{T} K^2 \left(r_N + K\right)^2} \cdot \frac{\mu\, e^{-2T}}{4\sqrt{T}} \cdot 2^{\left(\alpha^{(N-1)}\right)}
\end{aligned} \tag{61}$$

for every $N \in \mathbb{N}$. This shows

$$\begin{aligned}
\lim_{N \to \infty} \mathbb{E}\left|Y_N^N\right| &\geq \lim_{N \to \infty} \left(\mathbb{P}\left[\Omega_N\right] \cdot 2^{\left(\alpha^{(N-1)}\right)}\right) \\
&\geq \frac{\mu\, e^{-2T}}{4\sqrt{T}} \cdot \lim_{N \to \infty} \left(\left(\frac{1}{2}\sqrt{\frac{T}{N}}\right)^{N} \cdot e^{-\frac{N}{T} K^2 \left(r_N + K\right)^2} \cdot 2^{\left(\alpha^{(N-1)}\right)}\right) = \infty.
\end{aligned}$$

Due to Jensen's inequality we conclude

$$\lim_{N \to \infty} \mathbb{E}\left|Y_N^N\right|^p = \infty. \tag{62}$$

Hence, since $\mathbb{E}\left|X_T\right|^p < \infty$, we have

$$\lim_{N\to\infty} \left|\mathbb{E}\left|X_T\right|^p - \mathbb{E}\left|Y_N^N\right|^p\right| = \infty, \qquad \lim_{N\to\infty} \mathbb{E}\left|X_T - Y_N^N\right|^p = \infty, \tag{63}$$

which is the assertion. □

References

[1] Berkaoui, A., Bossy, M., and Diop, A. Euler scheme for SDEs with non-Lipschitz diffusion coefficient: strong convergence. *ESAIM Probab. Stat. 12* (2008), 1–11 (electronic).

[2] Bernard, P., and Fleury, G. Convergence of numerical schemes for stochastic differential equations. *Monte Carlo Methods Appl. 7*, 1-2 (2001), 35–44. Monte Carlo and probabilistic methods for partial differential equations (Monte Carlo, 2000).

[3] Deuflhard, P., and Bornemann, F. *Scientific computing with ordinary differential equations*, vol. 42 of *Texts in Applied Mathematics.* Springer-Verlag, New York, 2002. Translated from the 1994 German original by Werner C. Rheinboldt.

[4] Fleury, G. Convergence of schemes for stochastic differential equations. *Prob. Eng. Mech. 21* (2005), 35–43.

[5] Ginzburg, V. L., and Landau, L. D. *J. Exptl. Theoret. Phys. (U.S.S.R) 20*, 1064 (1950).

[6] Gyöngy, I. A note on Euler's approximations. *Potential Anal. 8*, 3 (1998), 205–216.

[7] Higham, D., and Mao, X. Convergence of Monte Carlo Simulations involving the Mean-Reverting Square Root Process. *Journal of Computational Finance, 8*, 3 (2005), 35–62.

[8] Higham, D. J., Mao, X., and Stuart, A. M. Strong convergence of Euler-type methods for nonlinear stochastic differential equations. *SIAM J. Numer. Anal. 40*, 3 (2002), 1041–1063 (electronic).

[9] Hu, Y. Semi-implicit Euler-Maruyama scheme for stiff stochastic equations. In *Stochastic analysis and related topics, V (Silivri, 1994)*, vol. 38 of *Progr. Probab.* Birkhäuser Boston, Boston, MA, 1996, pp. 183–202.

[10] Hutzenthaler, M., and Wakolbinger, A. Ergodic behavior of locally regulated branching populations. *Ann. Appl. Probab. 17*, 2 (2007), 474–501.

[11] Jentzen, A., and Kloeden, P. E. Pathwise taylor schemes for random ordinary differential equations. *BIT Numer. Math. 49* (2009), 113–140.

[12] Jentzen, A., Kloeden, P. E., and Neuenkirch, A. Pathwise approximation of stochastic differential equations on domains: higher order convergence rates without global lipschitz coefficients. *Numer. Math. 112* (2009), 41–64.

[13] Klenke, A. *Probability theory.* Universitext. Springer-Verlag London Ltd., London, 2008. A comprehensive course, Translated from the 2006 German original.

[14] Kloeden, P. E., and Platen, E. *Numerical solution of stochastic differential equations*, vol. 23 of *Applications of Mathematics (New York).* Springer-Verlag, Berlin, 1992.

[15] Lamba, H., Mattingly, J. C., and Stuart, A. M. An adaptive Euler-Maruyama scheme for SDEs: convergence and stability. *IMA J. Numer. Anal. 27*, 3 (2007), 479–506.

[16] Lambert, A. The branching process with logistic growth. *Ann. Appl. Probab. 15*, 2 (2005), 1506–1535.

[17] Lemaire, V. An adaptive scheme for the approximation of dissipative systems. *Stochastic Process. Appl. 117*, 10 (2007), 1491–1518.

[18] Mao, X. *Stochastic differential equations and their applications.* Horwood Publishing Series in Mathematics & Applications. Horwood Publishing Limited, Chichester, 1997.

[19] Mattingly, J. C., Stuart, A. M., and Higham, D. J. Ergodicity for SDEs and approximations: locally Lipschitz vector fields and degenerate noise. *Stochastic Process. Appl. 101*, 2 (2002), 185–232.

[20] Milstein, G. N. *Numerical integration of stochastic differential equations*, vol. 313 of *Mathematics and its Applications*. Kluwer Academic Publishers Group, Dordrecht, 1995. Translated and revised from the 1988 Russian original.

[21] Milstein, G. N., and Tretyakov, M. V. Numerical integration of stochastic differential equations with nonglobally Lipschitz coefficients. *SIAM J. Numer. Anal. 43*, 3 (2005), 1139–1154 (electronic).

[22] Ohta, T., and Kimura, M. A model of mutation appropriate to estimate the number of electrophoretically detectable alleles in a finite population. *Genet. Res. 22*, 2 (1973), 201–204.

[23] Talay, D. Stochastic Hamiltonian systems: exponential convergence to the invariant measure, and discretization by the implicit Euler scheme. *Markov Process. Related Fields 8*, 2 (2002), 163–198. Inhomogeneous random systems (Cergy-Pontoise, 2001).

[24] Yan, L. The Euler scheme with irregular coefficients. *Ann. Probab. 30*, 3 (2002), 1172–1194.

[25] Yuan, C., and Mao, X. A note on the rate of convergence of the Euler-Maruyama method for stochastic differential equations. *Stoch. Anal. Appl. 26*, 2 (2008), 325–333.

[26] Zhang, X. Euler-Maruyama approximations for SDEs with non-Lipschitz coefficients and applications. *J. Math. Anal. Appl. 316*, 2 (2006), 447–458.